

Quality ID #223 (NQF 0428): Functional Status Change for Patients with General Orthopaedic Impairments – National Quality Strategy Domain: Communication and Care Coordination

2018 OPTIONS FOR INDIVIDUAL MEASURES:

REGISTRY ONLY

MEASURE TYPE:

Outcome

DESCRIPTION:

A self-report outcome measure of functional status (FS) for patients 14 years+ with general orthopaedic impairments (neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment). The change in FS assessed using FOTO (general orthopedic) PROM (patient reported outcomes measure) is adjusted to patient characteristics known to be associated with FS outcomes (risk adjusted) and used as a performance measure at the patient level, at the individual clinician, and at the clinic level by to assess quality

INSTRUCTIONS:

This outcomes measure is to be submitted **once per treatment episode** for all patients with a functional deficit related to the neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment. This is an outcomes measure and its calculation requires submitting of the patient's functional status score, as a minimum, at admission to and again at discharge from an episode of rehabilitation. The admission score, estimated using patient self-report surveys, is recorded during the first rehabilitation treatment encounter and the discharge score is recorded at or near the conclusion of the final rehabilitation treatment encounter. It is anticipated that physical and occupational therapists providing treatment for functional neck, cranium, mandible, thoracic spine, ribs or other general orthopedic deficits will submit this measure.

Definitions:

Functional deficit – Limitation or impairment of physical abilities/function resulting in evaluation and inclusion in a treatment plan of care.

Treatment Episode – A Treatment Episode is defined as beginning with an Admission for a functional neck, cranium, mandible, thoracic spine, ribs or other general orthopedic deficit, progressing to development of a plan of care, including treatment, without interruption of care (for example, a hospitalization or surgical intervention), and ending with Discharge from clinical care by the eligible clinician. A patient currently under clinical care for a neck, cranium, mandible, thoracic spine, ribs or other general orthopedic deficit remains in a single episode of care until the Discharge is conducted and documented by the eligible clinician.

Admission – An Admission is the first encounter for a functional deficit involving the neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment and includes an evaluation (CPT 97161, 97162, 97163 for physical therapy or 97165, 97166, 97167 for occupational therapy) and development of a plan of care by the eligible clinician. A patient presenting with a neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment, who has had an interruption of a Treatment Episode for the same functional neck, cranium, mandible, thoracic spine, ribs or other general orthopedic deficit secondary to an appropriate reason like hospitalization or surgical intervention, is a new Admission.

Discharge – Discharge is accompanied by a re-evaluation (CPT 97164 for physical therapy or 97168 for occupational therapy) or Functional Limitation Submitting Discharge Status G-Code (G8980, G8983, G8986, G8989, G8992 or G8995) identifying the close of a Treatment Episode for the same neck, cranium, mandible, thoracic spine, ribs or other general orthopedic deficit identified at admission and documented by a discharge report by the eligible clinician. An interruption in clinical care for an appropriate reason like hospitalization or surgical intervention requires a discharge from the current Treatment Episode.

Encounter – A face to face visit between the patient and the provider for the purpose of assessing and/or improving a functional deficit.

Patient Reported – The patient directly, or through a proxy, provides answers to functional status survey items using standardized, reliable and valid, computerized adaptive testing or paper and pencil survey methods.

Measure Submission:

The listed denominator criteria is used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions allowed by the measure. The quality-data codes listed do not need to be submitted for registry submissions; however, these codes may be submitted for those registries that utilize claims data.

DENOMINATOR:

All patients 14 years and older with general orthopedic impairments (neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment) who have initiated rehabilitation treatment and completed the FOTO (general orthopaedic) PROM

Option 1 – Physical Therapy Denominator Criteria (Eligible Cases):

All patients aged ≥ 14 years on date of encounter

AND

Patient encounter during the performance period identifying evaluation (CPT): 97161, 97162, 97163

AND

Patient encounter during the performance period identifying discharge (CPT or HCPCS): 97164, G8980, G8983, G8986, G8989, G8992, G8995

AND

Functional deficit affecting neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment

AND NOT

DENOMINATOR EXCLUSIONS:

Patient refused to participate: G9738

OR

Patient unable to complete the FOTO General Orthopedic Intake PROM at admission and discharge due to blindness, illiteracy, severe mental incapacity or language incompatibility and an adequate proxy is not available: G9739

OR

Option 2 – Occupational Therapy Denominator Criteria (Eligible Cases):

All patients aged ≥ 14 years on date of encounter

AND

Patient encounter during the performance period identifying evaluation (CPT): 97165, 97166, 97167

AND

Patient encounter during the performance period identifying discharge (CPT or HCPCS): 97168, G8980, G8983, G8986, G8989, G8992, G8995

AND

Functional deficit affecting neck, cranium, mandible, thoracic spine, ribs or other general orthopedic impairment

AND NOT

DENOMINATOR EXCLUSIONS:

Patient refused to participate: G9738

OR

Patient unable to complete the FOTO General Orthopedic Intake PROM at admission and discharge due to blindness, illiteracy, severe mental incapacity or language incompatibility and an adequate proxy is not available: G9739

NUMERATOR:

Patients who were presented with the FOTO General Functional Status measure at Admission (Intake) and Discharge (Status) for the purpose of calculating the patient's Risk-adjusted Functional Status Change Residual Score

Definitions:

Patient's Functional Status Score – A functional status score is produced when the patient completes the FOTO functional status survey (either by paper and pencil or computerized adaptive testing administration). The functional status score is continuous and linear. Scores range from 0 to 100 with higher scores meaning higher functional abilities. The survey is standardized, and the scores are validated for the measurement of function for this population

Patient's Functional Status Change Score – A functional status change score is calculated by subtracting the Patient's Functional Status Score at Admission from the Patient's Functional Status Score at Discharge.

Predicted Functional Status Change Score – Functional Status Change Scores for patients are risk adjusted using multiple linear regression methods that include the following independent variables: Patient's Functional Status Score at Admission, patient age, symptom acuity, surgical history, gender, number of co-morbidities, and level of fear-avoidance. The Patient's Functional Status Change Score is the dependent variable. The statistical regression produces a Risk-Adjusted Predicted Functional Status Change Score.

Risk-Adjusted Functional Status Change Residual Score – The difference between the raw non-risk-adjusted Patient's Functional Status Change Score and the Risk-Adjusted Predicted Functional Status Change Score (raw minus predicted) is the Risk-Adjusted Functional Status Change Residual Score, which is in the same units as the Patient's Functional Status Scores, and should be interpreted as the unit of functional status change different than predicted given the risk-adjustment variables of the patient being treated. As such, the Risk-Adjusted Residual Change Score represents Risk-Adjusted Change corrected for the level of severity of the patient. Risk-Adjusted Residual Change Scores of zero (0) or greater (> 0) should be interpreted as functional status change scores that were predicted or better than predicted given the risk-adjustment variables of the patient, and risk-adjusted residual change scores less than zero (< 0) should be interpreted as functional status change scores that were less than predicted given the risk-adjustment variables of the patient. Aggregated Risk-Adjusted Residual Scores allow meaningful comparisons amongst clinicians or clinics.

Not Appropriate (Denominator Exception) – Prior to conclusion of Plan of Care, intervention was interrupted or discontinued for any reason including by the referring physician, the provider, the payer or the patient, and attempts by the provider to complete a follow-up functional status survey near Discharge were unsuccessful.

Numerator Options:

Performance Met:

Risk-Adjusted Functional Status Change Residual Score for the neck, cranium, mandible, thoracic spine, ribs or other general orthopaedic impairment successfully calculated and the score was equal to zero (0) or greater than zero (> 0) (G8671)

OR

Performance Met:

Risk-Adjusted Functional Status Change Residual Score for the neck, cranium, mandible, thoracic spine, ribs or other general orthopaedic impairment successfully calculated and the score was less than zero (< 0) (G8672)

OR

Denominator Exception:

Risk-Adjusted Functional Status Change Residual Scores for the neck, cranium, mandible, thoracic spine, ribs or other general orthopaedic impairment not measured because the patient did not complete FOTO's Functional follow up Status Survey near discharge, patient Not Appropriate (G8673)

OR

Performance Not Met:

Risk-Adjusted Functional Status Change Residual Scores for the neck, cranium, mandible, thoracic spine, ribs or other general orthopaedic impairment not measured because the patient did not complete FOTO's Functional Intake on admission and/or follow up Status Survey near discharge, reason not given (G8674)

RATIONALE:

Functional deficits are common in the general population and are costly to the individual, their family and society. Improved functional status has been associated with greater quality of life, self-efficacy, improved financial well-being and lower future medical costs. Improving functional status in people seeking rehabilitation has become a goal of the

American Physical Therapy Association. Therefore, measuring change in functional status is important for providers treating patients in rehabilitation and can be used to assess the success of treatment and direct modification of treatment.

Change in functional status represents the activity domain of the International Classification of Function. If treatment is designed to improve the functional deficit, it is logical to assess functional status at discharge using a standardized score to determine if treatment improved the functional status of the patient over the treatment episode.

The National Quality Measures Clearinghouse has approved the measurement of change in functional status, using this survey. (NQMC-0022)

CLINICAL RECOMMENDATION STATEMENTS:

The American Physical Therapy Association (APTA), in their Guide to Physical Therapy Practice, described five recommended elements of patient management: examination, evaluation, diagnosis, prognosis and intervention. The elements were intended to direct therapists in their approach to patient treatment for the purpose of optimizing patient outcomes. The APTA clearly identifies functional status data as one of the major forms of data to be collected for patients receiving rehabilitation. The functional status measures should be used to assist in the planning, implementation and modification of treatment interventions and should be used as measures of outcomes. The current functional status scores can be used by therapists to fulfill the recommended methods of the APTA in the management of patients in rehabilitation.

COPYRIGHT:

Functional Health Status (FS) Measures and related data specifications have been developed by Focus On Therapeutic Outcomes, Inc. (FOTOTM). The measures are calculated from data gathered by patient self-report surveys that are copyrighted by FOTOTM and exist in two formats: electronic and short form paper formats.

Proprietary computerized adaptive testing electronic surveys and risk adjusted FS measure reports are included in Patient Inquiry®, a web based survey engine published by FOTOTM. The short-form surveys and the accompanying scoring and risk-adjustment specifications, while copyrighted, can be downloaded, at:

Link to access all Measures: [Link to All FOTO Measures](#)

Therefore, the short-form surveys and accompanying scoring and risk-adjustment specifications may be reproduced and distributed, without modification, for noncommercial purposes, e.g., **use by health care providers in connection with their practices**. Commercial use is defined as the sale, license, or distribution of the surveys or Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Measures require a license agreement between the user and Focus On Therapeutic Outcomes, Inc.

Limited proprietary coding is contained in the Measure specifications for convenience.


THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND.

© 2015 Focus On Therapeutic Outcomes Inc. All Rights Reserve

FOTO® disclaims all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

CPT® contained in the Measures specifications is copyright 2004-2017 American Medical Association.

**2018 Registry Flow for Quality ID #223 NQF #0428:
Functional Status Change for Patients with General Orthopaedic Impairments**



* See the posted Measure Specification for specific coding and instructions to submit this measure.
NOTE: Submission Frequency: Episode

CPT only copyright 2017 American Medical Association. All rights reserved.
The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

**2018 Registry Flow for Quality ID #223 NQF #0428:
Functional Status Change for Patients with General Orthopaedic Impairments**

SAMPLE CALCULATIONS:

Data Completeness=

$$\frac{\text{Performance Met (a}^1+\text{a}^2=30 \text{ episodes)} + \text{Denominator Exception (b=20 episodes)} + \text{Performance Not Met (c=20 episodes)}}{\text{Eligible Population / Denominator (d=80 episodes)}} = \frac{70 \text{ episodes}}{80 \text{ episodes}} = 87.50\%$$

Performance Rate=

$$\frac{\text{Performance Met (a}^1+\text{a}^2=30 \text{ episodes)}}{\text{Data Completeness Numerator (70 episodes) - Denominator Exception (b=20 episodes)}} = \frac{30 \text{ episodes}}{50 \text{ episodes}} = 60.00\%$$

* See the posted Measure Specification for specific coding and instructions to submit this measure.

NOTE: Submission Frequency: Episode

CPT only copyright 2017 American Medical Association. All rights reserved.
The measure diagrams were developed by CMS as a supplemental resource to be used in conjunction with the measure specifications. They should not be used alone or as a substitution for the measure specification.

v2

2018 Registry Flow for Quality ID

#223 NQF #0428: Functional Status Change for Patients with General Orthopedic Impairments

Please refer to the specific section of the specification to identify the denominator and numerator information for use in submitting this Individual Specification. This flow is for registry data submission..

1. Start with Denominator
2. Check Patient Age:
 - a. If Age greater than or equal to 14 years of age on Date of Service equals No during the measurement period, do not include in Eligible Patient Population. Stop Processing.
 - b. If Age greater than or equal to 14 years of age on Date of Service equals Yes during the measurement period, proceed to check Patient Evaluation Encounter During Performance period .
3. Check Physical Therapy (PT) Patient Evaluation Encounter During Performance period:
 - a. If Patient PT Evaluation Encounter During Performance period equals No, proceed to Patient Occupational Therapy (OT) Encounter Evaluation during the Performance period.
 - b. If Patient PT Evaluation Encounter During Performance period equals Yes, proceed to Patient (PT) Encounter During the Performance period Identifying Discharge.
4. Check Patient (PT) Encounter During the Performance period Identifying Discharge:
 - a. If Patient PT Encounter During the Performance period Identifying Discharge and as listed in the denominator equals No, proceed to Patient OT Encounter Evaluation during the Performance period.
 - b. If Patient PT Encounter During the Performance period Identifying Discharge and as listed in the denominator equals Yes, proceed to Functional Deficit Affecting General Orthopedic Impairments.
5. Check Patient OT Encounter Evaluation during the Performance period:
 - a. If Patient OT Encounter Evaluation during the Performance period and as listed in the Denominator equals No, do not include in Eligible Population. Stop Processing.
 - b. If Patient OT Encounter Evaluation during the Performance period and as listed in the Denominator equals Yes, proceed to Patient OT Encounter During Performance period Identifying Discharge.
6. Check Patient OT Encounter During Performance period Identifying Discharge:
 - a. If Patient OT Encounter During Performance period Identifying Discharge as listed in the Denominator equals Yes, proceed to Functional Deficit Affecting General Orthopedic Impairments.
7. Check Functional Deficit Affecting General Orthopedic Impairments:
 - a. If Functional Deficit Affecting General Orthopedic Impairments equals No, do not include in Eligible Patient Population. Stop Processing.
 - b. If Functional Deficit Affecting General Orthopedic Impairments equals Yes, proceed to Denominator Exclusion, Patient Refused to Participate.
8. Check Denominator Exclusion, Patient Refused to Participate:

- a. If Denominator Exclusion, Patient Refused to Participate equals Yes, do not include in Eligible Population. Stop Processing.
 - b. If Denominator Exclusion, Patient Refused to Participate equals No, proceed to check next Denominator Exclusion, Patient Unable to Complete the FOTO General Orthopedic Intake PROM at Admission and Discharge.
9. Check Denominator Exclusion, Patient Unable to Complete the FOTO General Orthopedic Intake PROM at Admission and Discharge:
 - a. If Denominator Exclusion, Patient Unable to Complete the FOTO General Orthopedic Intake PROM at Admission and Discharge equals Yes, do not include in Eligible Population. Stop Processing.
 - b. If Denominator Exclusion, Patient Unable to Complete the FOTO General Orthopedic Intake PROM at Admission and Discharge equals No, include in Eligible Population and proceed to Denominator Population.
10. Denominator Population:
 - a. Denominator population is all Eligible Patients in the Denominator. Denominator is represented as Denominator in the Sample Calculation listed at the end of this document. Letter d equals 80 episodes in the Sample Calculation.
11. Start Numerator
12. Check Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was greater than or equal to 0:
 - a. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was greater than or equal to 0 equals Yes, include in Data Completeness Met and Performance Met.
 - b. Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a1 equals 10 episodes in the Sample Calculation.
 - c. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was greater than or equal to 0 equals No, proceed to Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was less than 0.
13. Check Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was less than 0:
 - a. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was less than 0 equals Yes, include in Data Completeness Met and Performance met.
 - b. Data Completeness Met and Performance Met letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter a2 equals 20 episodes in the Sample Calculation.

- c. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Successfully Calculated and the Score was less than 0 equals No, proceed to Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured, Patient Not Eligible/Not Appropriate.
14. Check Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured, Patient Not Eligible/Not Appropriate
- a. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured, Patient Not Eligible/Not Appropriate equals Yes, include in the Data Completeness Met and Denominator Exception.
 - b. Data Completeness Met and Denominator Exception letter is represented in the Data Completeness and Performance Rate in the Sample Calculation listed at the end of this document. Letter b equals 20 episodes in the Sample Calculation.
 - c. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured, Patient Not Eligible/Not Appropriate equals No, proceed to Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured, because the patient did not complete FOTO's Functional Intake on admission and/or Follow Up Status Survey near discharge Reason Not Given.
15. Check Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured because the patient did not complete FOTO's Functional Intake on admission and/or Follow Up Status Survey near discharge Reason Not Given:
- a. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured because the patient did not complete FOTO's Functional Intake on admission and/or Follow Up Status Survey near discharge, Reason Not Given equals Yes, include in the Data Completeness Met and Performance Not Met.
 - b. Data Completeness Met and Performance Not Met letter is represented in the Data Completeness in the Sample Calculation listed at the end of this document. Letter c equals 20 episodes in the Sample Calculation.
 - c. If Risk-Adjusted Functional Status Change Residual Score for the Neck, Cranium, Mandible, Thoracic Spine, Ribs, or Other General Orthopedic Impairments Not Measured because the patient did not complete FOTO's Functional Intake on admission and/or Follow Up Status Survey near discharge, Reason Not Given equals No, proceed to Data Completeness Not Met.
16. Check Data Completeness Not Met
- a. If Data Completeness Not Met equals No, Quality Data Code or equivalent not submitted. 10 episodes have been subtracted from Data Completeness Numerator in the Sample Calculation.

SAMPLE CALCULATIONS:

Data Completeness=

$$\frac{\text{Performance Met (a}^1+\text{a}^2=30 \text{ episodes)} + \text{Denominator Exception (b=20 episodes)} + \text{Performance Not Met (c=20 episodes)}}{\text{Eligible Population / Denominator (d=80 episodes)}} = \frac{70 \text{ episodes}}{80 \text{ episodes}} = 87.50\%$$

Performance Rate=

$$\frac{\text{Performance Met (a}^1+\text{a}^2=30 \text{ episodes)}}{\text{Data Completeness Numerator (70 episodes) - Denominator Exception (b=20 episodes)}} = \frac{30 \text{ episodes}}{50 \text{ episodes}} = 60.00\%$$